

CT Coronary Angiography for Evaluation of Subclinical Atherosclerosis in Patients with High-Risk Metabolic Syndrome

Dr Divya Kumari¹, Dr. Dilip Kumar²

1. Department of Radiodiagnosis, SK Medical College and Hospital, UP
2. Department of General Medicine, SK Medical College and Hospital, UP

Received: 30-04-2026 / Accepted: 15-05-2026/ Published:28-05-2026

Corresponding author: Dr Divya Kumari

Conflict of interest: Nil

ABSTRACT

Background:

Metabolic syndrome (MetS) is a major driver of premature atherosclerotic cardiovascular disease, particularly in South Asian populations. Conventional risk scores often fail to detect early or subclinical coronary disease in high-risk individuals. Coronary CT angiography (CCTA) enables direct visualisation and quantification of coronary plaque, including high-risk morphology. This study evaluates the role of CCTA in detecting subclinical atherosclerosis among adults with high-risk MetS.

Methods:

This cross-sectional observational study included 100 adults diagnosed with MetS (NCEP-ATP III criteria) and classified as high-risk based on ≥ 4 MetS components, long-standing MetS, or additional atherosclerotic risk enhancers. All participants underwent contrast-enhanced CCTA on a ≥ 64 -slice CT scanner. Plaque presence, type, burden (segment involvement score, SIS), stenosis severity, high-risk plaque features, and coronary artery calcium score (CACS) were assessed. Metabolic parameters were correlated with plaque burden.

Results:

Subclinical coronary atherosclerosis was detected in 72% of patients. Plaque prevalence rose significantly with MetS severity (40.9% in mild vs 92.5% in severe; $p < 0.001$). Calcified (22.7% vs 60%; $p = 0.004$) and non-calcified plaques (9.1% vs 42.5%; $p = 0.008$) increased across severity categories. High-risk plaque features were present in 32.5% of severe MetS subjects compared with 4.5% in mild ($p = 0.003$). Median CACS (42 to 152; $p < 0.001$) and mean SIS (0.9 ± 0.7 to 3.0 ± 1.4 ; $p < 0.001$) also increased markedly. Luminal stenosis $\geq 50\%$ occurred in 40% of severe MetS patients but was absent in mild cases. Plaque burden showed significant correlations with fasting glucose ($r = 0.42$), triglycerides ($r = 0.38$), waist circumference ($r = 0.36$), BMI ($r = 0.29$), and HDL ($r = -0.31$). MetS severity score demonstrated the strongest association ($\rho = 0.55$; $p < 0.001$).

Conclusion:

Subclinical coronary atherosclerosis is highly prevalent in high-risk MetS patients, with plaque burden and high-risk features rising sharply with metabolic severity. CCTA offers substantial value in early detection and risk stratification, surpassing traditional clinical and biochemical markers. Incorporating CCTA into evaluation algorithms for high-risk MetS populations may facilitate earlier preventive interventions. Prospective studies are required to determine the impact of CCTA-guided management on long-term cardiovascular outcomes.

Keywords: Metabolic syndrome, CT coronary angiography, subclinical atherosclerosis, coronary plaque, high-risk plaque, coronary calcium score.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Metabolic syndrome (MetS)—characterised by central obesity, insulin resistance, dyslipidaemia, hypertension and a pro-inflammatory/pro-thrombotic state—has emerged as a major global driver of atherosclerotic cardiovascular disease (ASCVD) (1–3). Individuals with MetS have a substantially elevated risk of overt coronary artery disease (CAD) and myocardial infarction compared

with those without MetS, even after adjustment for conventional cardiovascular risk factors (4). The atherosclerotic process in MetS remains silent and subclinical for many years before progressing to symptomatic disease (5). Traditional cardiovascular risk prediction tools, such as the Framingham Risk Score and SCORE system offer useful population-level estimates but are limited in detecting early or

subclinical atherosclerosis in high-risk individuals (6–7). Consequently, non-invasive imaging modalities have gained prominence for identifying subclinical coronary disease before the onset of symptoms, thereby enabling improved patient selection for early preventive therapy (7–9).

Among available imaging options, computed tomography (CT)–based techniques have become central to the non-invasive evaluation of coronary atherosclerosis. Coronary CT angiography (CCTA), in particular, allows direct in vivo visualization and quantification of coronary plaque—including calcified, non-calcified, and mixed components—and provides detailed assessments of plaque burden, luminal stenosis, positive remodelling, and high-risk plaque characteristics (10–12). These capabilities extend beyond coronary artery calcium (CAC) scoring alone, which, while useful in risk prediction, reflects only the calcified component of atherosclerosis and may underestimate early disease (13–15).

Evidence from asymptomatic and intermediate-risk populations suggests that even non-obstructive plaque detected by CCTA is significantly associated with future cardiovascular events, independent of traditional risk factors (16–17). Thus, CCTA offers a more sensitive method for early detection of subclinical coronary atherosclerosis and may refine risk stratification in high-risk individuals.

In MetS, the pro-atherogenic environment created by insulin resistance, dyslipidaemia, endothelial dysfunction, and chronic low-grade inflammation accelerates the progression of coronary plaque, even in the absence of symptoms (18–19). While previous research has demonstrated associations between MetS components and surrogate markers such as carotid intima–media thickness and coronary calcium score (7,13), there remains a relative paucity of data specifically evaluating the utility of CCTA in characterising subclinical coronary atherosclerosis in MetS patients (20–21).

Given the rising prevalence of MetS in South Asia and other developing regions, coupled with the significant burden of premature CAD (1–3,22), early detection of subclinical coronary atherosclerosis is crucial for guiding individualised preventive strategies. CCTA may offer significant clinical value by identifying silent coronary plaque, quantifying total plaque burden, and revealing high-risk plaque morphology in patients with MetS who might otherwise be underestimated by conventional risk scores. Therefore, the present study aims to evaluate the role of CT coronary angiography in detecting and characterising subclinical atherosclerosis in adults with high-risk metabolic syndrome.

Methodology:

This was a hospital-based cross-sectional observational study conducted in the Department of Radiology and medicine, Department of medicine, at a tertiary care teaching hospital. A total of 100 consecutive adults with metabolic syndrome (MetS) who fulfilled the inclusion criteria were enrolled.

Study Design

The present study was a cross-sectional observational study conducted to evaluate cardiac dysfunction by echocardiography in patients with Chronic Obstructive Pulmonary Disease (COPD). Patients were classified according to the Global Initiative for Chronic Obstructive Lung Disease (GOLD) Guideline 2024 based on spirometry findings.

Study Setting

The study was conducted in the Departments of Respiratory Medicine and General Medicine at K.D. Medical College Hospital and Research Centre, Mathura. The hospital catered to both urban and rural populations, ensuring inclusion of a diverse group of COPD patients.

Eligibility Criteria

Inclusion Criteria

1. Adults aged ≥ 18 years diagnosed with Metabolic Syndrome as per NCEP-ATP III criteria (presence of ≥ 3 of the following):

- Abdominal obesity
- Elevated fasting blood glucose
- Elevated triglycerides
- Reduced HDL-cholesterol
- Elevated blood pressure

2. Patients categorized as high-risk, defined as:

- Presence of ≥ 4 MetS components, or
- Long-standing MetS (≥ 5 years), or
- Additional atherosclerotic risk enhancers (smoking, family history of premature CAD, chronic inflammatory state)

3. Patients who underwent contrast-enhanced coronary CT angiography (CCTA) during the study period.

Exclusion Criteria

- Known coronary artery disease (prior MI, PCI, CABG).
- Symptomatic patients with typical angina.
- Renal impairment (eGFR < 60 mL/min/1.73 m²).
- Contrast allergy or contraindication to iodinated contrast.
- Pregnancy or lactation.

- Inadequate CT image quality (motion artefact, arrhythmia).

Clinical and Laboratory Assessment

All participants underwent a detailed clinical evaluation including:

- Age, sex, anthropometry (BMI, waist circumference)
- Blood pressure measurement
- Lifestyle factors (smoking, alcohol, activity levels)
- Biochemical parameters performed after overnight fasting included:
 - Fasting blood glucose
 - Lipid profile—LDL, HDL, triglycerides
 - HbA1c
 - Renal and liver function tests

MetS severity categories were documented based on number of components and risk enhancers.

CT Coronary Angiography Protocol

Scanner and Acquisition

CCTA was performed using a 64-slice or higher multi-detector CT scanner.

Scan parameters included:

- Retrospective or prospective ECG-gated acquisition
- Tube voltage: 100–120 kVp
- Tube current modulation applied
- Slice thickness: 0.6–0.75 mm
- Reconstruction in axial, MPR, curved-MPR, and 3D volume-rendered formats

Contrast Administration

- 60–80 mL of non-ionic iodinated contrast injected at 4–5 mL/s
- Bolus tracking used to initiate acquisition
- Beta-blockers administered when required to achieve HR <65 bpm
- Sublingual nitroglycerin given unless contraindicated

Image Analysis

All images were evaluated independently by two experienced radiologists blinded to clinical details. Discrepancies were resolved by consensus.

Parameters Assessed

1. **Presence of coronary plaque (yes/no).**
2. **Plaque characterization:**
 - Calcified
 - Non-calcified
 - Mixed

3. **Segment involvement score (SIS) – number of coronary segments with plaque.**

4. **Severity of luminal stenosis:**
 - <25%
 - 25–49%
 - 50–69%
 - ≥70%

5. **High-risk plaque features:**
 - Positive remodelling
 - Low-attenuation plaque
 - Napkin-ring sign
 - Spotty calcification

6. **Coronary Artery Calcium Score (CACS) measured using Agatston method.**

Outcome Measures

Primary outcome:

- Prevalence and extent of subclinical coronary atherosclerosis in MetS.

Secondary outcomes:

- Association between MetS severity and:
 - Plaque presence
 - Plaque type
 - Plaque burden
 - High-risk features
 - CACS and stenosis severity

Statistical Analysis

Data were analysed using SPSS version 25.

- **Categorical variables:** expressed as frequencies and percentages.
- Continuous variables: mean ± SD or median (IQR).
- Group comparisons performed using:
 - Chi-square / Fisher's exact test for categorical variables
 - Independent t-test / ANOVA for normally distributed variables
 - Mann–Whitney U / Kruskal–Wallis test for skewed variables
 - Correlation analysis using Pearson or Spearman coefficients.
 - A p-value < 0.05 was considered statistically significant.

Results:

The demographic characteristics of the study participants, as shown in Table 1, demonstrate that the mean age was 52.6 ± 9.4 years, with a median of 53 years, indicating a predominantly middle-aged population. Age increased significantly with rising

metabolic syndrome severity ($p = 0.004$), suggesting that advancing age may contribute to worsening metabolic derangement. Waist circumference and BMI showed similar significant upward trends across severity categories ($p = 0.001$ and $p = 0.010$, respectively), reinforcing the strong influence of central adiposity on disease progression. Table 2 presents the sex distribution, showing that although males constituted 58% of the cohort, no statistically significant relationship existed between sex and metabolic syndrome severity ($p = 0.63$), indicating that the severity of metabolic syndrome was not gender-dependent in this population. Hypertension was the most common comorbidity, present in 41% of patients, followed by diabetes mellitus in 25%. Hypothyroidism and cerebrovascular accident were observed in 5% and 4% of patients, respectively, while 25% of patients had no associated comorbidity. Smoking was highly prevalent, with 68% of patients being smokers and 32% non-smokers.

The biochemical and metabolic parameters summarised in Table 3 reveal progressively worsening abnormalities with increasing metabolic syndrome severity. Fasting glucose and triglyceride levels rose significantly ($p < 0.001$ and $p = 0.002$), HDL levels declined markedly ($p = 0.001$), and systolic blood pressure increased steadily ($p = 0.003$), reflecting the clustering and intensification of metabolic disturbances as the syndrome becomes more severe. Table 4 highlights the coronary CT angiography findings, where 72% of all participants exhibited subclinical coronary atherosclerosis despite being asymptomatic. A striking gradient was observed, with plaque prevalence rising from 40.9% in mild cases to 92.5% in severe metabolic syndrome ($p < 0.001$). Both calcified and non-

calcified plaques demonstrated significant increases with severity ($p = 0.004$ and $p = 0.008$), while high-risk plaque features were especially notable in severe metabolic syndrome, occurring in 32.5% of this group compared with only 4.5% of mild cases ($p = 0.003$). The coronary artery calcium score and the segment involvement score also increased markedly with severity (both $p < 0.001$), indicating not only a higher frequency but a greater burden of coronary disease. Additionally, luminal stenosis $\geq 50\%$ was absent in the mild group but present in 40% of severe metabolic syndrome subjects ($p < 0.001$), underscoring the silent progression of potentially obstructive coronary disease.

Table 5 further illustrates the relationship between metabolic parameters and plaque burden, showing significant positive correlations between plaque burden and fasting glucose ($r = 0.42$), triglycerides ($r = 0.38$), BMI ($r = 0.29$), and waist circumference ($r = 0.36$), while HDL exhibited a moderate negative correlation ($r = -0.31$). The strongest association was observed between the metabolic syndrome severity score and plaque burden ($p = 0.55$, $p < 0.001$), demonstrating that worsening metabolic derangement corresponded with increased coronary involvement. Collectively, the findings from Tables 1 through 5 clearly indicate that metabolic syndrome severity is strongly associated with both the presence and the extent of subclinical coronary atherosclerosis, highlighting CT coronary angiography as an effective tool for early detection and risk stratification in high-risk metabolic syndrome patients.

Table 1. Demographic Profile of Study Population (n = 100)

Variable	Mean \pm SD	Median (IQR)	Stage Comparison (MetS Severity)	Statistical Test	p-value
Age (years)	52.6 \pm 9.4	53 (46–59)	Mild (n=22): 48.3 \pm 7.5 Moderate (n=38): 52.4 \pm 9.1 Severe (n=40): 55.8 \pm 9.9	ANOVA	0.004
Waist Circumference (cm)	101.6 \pm 9.8	100 (95–108)	Mild: 96.2 \pm 8.7 Moderate: 101.3 \pm 9.1 Severe: 105.8 \pm 9.3	ANOVA	0.001
BMI (kg/m ²)	29.4 \pm 3.8	29.1 (27.0–31.4)	Mild: 27.6 \pm 3.1 Moderate: 29.1 \pm 3.4 Severe: 30.7 \pm 3.7	ANOVA	0.010

Table 2. Sex Distribution Across Metabolic Syndrome Severity Groups

Sex	Total (n=100)	Mild (n=22)	Moderate (n=38)	Severe (n=40)	Statistical Test	p-value
Male	58 (58%)	11 (50%)	23 (60.5%)	24 (60%)	Chi-square	0.63
Female	42 (42%)	11 (50%)	15 (39.5%)	16 (40%)	—	—

Table 3. Metabolic and Biochemical Parameters

Parameter	Mean ± SD	Median (IQR)	Mild	Moderate	Severe	Statistical Test	p-value
Fasting Glucose (mg/dL)	143.6 ± 32.1	138 (122–165)	124.8 ± 21.5	140.2 ± 28.7	159.3 ± 34.8	ANOVA	<0.001
Triglycerides (mg/dL)	209.4 ± 51.3	198 (174–236)	184.2 ± 41.5	204.9 ± 45.1	228.6 ± 56.3	ANOVA	0.002
HDL (mg/dL)	38.8 ± 6.4	38 (34–43)	42.3 ± 5.2	38.7 ± 6.1	36.9 ± 6.2	ANOVA	0.001
Systolic BP (mmHg)	142 ± 16	140 (130–152)	133.5 ± 12.1	141.4 ± 13.7	148.0 ± 15.5	ANOVA	0.003

Table 4. Coronary Atherosclerosis Findings on CCTA

CCTA Parameter	Total (n=100)	Mild (n=22)	Moderate (n=38)	Severe (n=40)	Statistical Test	p-value
Any Coronary Plaque (present)	72 (72%)	9 (40.9%)	26 (68.4%)	37 (92.5%)	Chi-square	<0.001
Calcified Plaque	44 (44%)	5 (22.7%)	15 (39.4%)	24 (60%)	Chi-square	0.004
Non-calcified (soft) Plaque	28 (28%)	2 (9.1%)	9 (23.7%)	17 (42.5%)	Chi-square	0.008
Mixed Plaque	26 (26%)	3 (13.6%)	10 (26.3%)	13 (32.5%)	Chi-square	0.128
High-Risk Plaque Features (any)	18 (18%)	1 (4.5%)	4 (10.5%)	13 (32.5%)	Chi-square	0.003
Coronary Artery Calcium Score (Agatston)	118 ± 64	95 (44–178)	Mild: 42 ± 25 Moderate: 112 ± 55 Severe: 152 ± 68	Kruskal–Wallis	<0.001	
Segment Involvement Score (SIS)	2.1 ± 1.4	0.9 ± 0.7	1.8 ± 1.0	3.0 ± 1.4	ANOVA	<0.001
Luminal Stenosis ≥50%	21 (21%)	0 (0%)	5 (13.1%)	16 (40%)	Fisher’s Exact	<0.001

Table 5. Correlation Between Metabolic Parameters and Coronary Atherosclerosis

Metabolic Parameter	Correlation with Plaque Burden (SIS)	p-value
---------------------	--------------------------------------	---------

Fasting Glucose	r = 0.42	<0.001
Triglycerides	r = 0.38	0.001
HDL	r = -0.31	0.003
BMI	r = 0.29	0.004
Waist Circumference	r = 0.36	0.001
MetS Severity Score	r = 0.55	<0.001

Discussion:

In the present cross-sectional study of 100 high-risk metabolic syndrome (MetS) patients, we found that 72% had detectable coronary plaque by CT coronary angiography (CCTA) despite being asymptomatic. Notably, plaque prevalence rose from 40.9% in the mild MetS group to 92.5% in the severe group ($p < 0.001$). Calcified plaques increased from 22.7% to 60% ($p = 0.004$), non-calcified plaques from 9.1% to 42.5% ($p = 0.008$), and high-risk plaque features were present in 32.5% of the severe group (versus 4.5% in the mild; $p = 0.003$). The coronary artery calcium score (CACS) and segment involvement score (SIS) also climbed significantly with MetS severity (median CACS from 42 to 152, $p < 0.001$; mean SIS 0.9 ± 0.7 to 3.0 ± 1.4 , $p < 0.001$). Metabolic parameters such as fasting glucose ($r = 0.42$), triglycerides ($r = 0.38$), waist circumference ($r = 0.36$), BMI ($r = 0.29$) and HDL ($r = -0.31$) correlated significantly with plaque burden; the MetS severity score had the strongest correlation ($\rho = 0.55$, $p < 0.001$).

These findings corroborate and extend prior literature. For instance, Park et al. examined 5,213 asymptomatic individuals undergoing CCTA and found that MetS (present in ~39% of the cohort) was an independent predictor of significant coronary artery disease (CAD) (OR ~1.99) and of events (RR ~1.67) during median follow-up of 28 months. (22) Their results echo our observation that MetS is associated with a high prevalence of silent coronary atherosclerosis; however, they reported plaque prevalence rather in terms of significant stenosis rather than burden metrics, and their population was much larger. In another study by Kim et al., MetS was shown to predict progression of CACS, coronary stenosis and vulnerable plaque on CCTA (HR ~1.45 for stenosis/plaques after adjustment) in 2,426 asymptomatic subjects. Our results add to these by quantifying multiple plaque types, burden scores (SIS), high-risk features, and by stratifying according to severity of MetS, thus revealing a clear dose-response relationship. (24)

The high prevalence of non-calcified plaque (42.5% in severe MetS) and high-risk features in our cohort is of particular significance, because non-calcified and positively remodeled plaques are known to carry higher risk of rupture and acute events than calcified, stable plaques. Prior smaller studies in morbid obesity found among BMI > 40 kg/m²

patients that majority of plaques on CCTA were non-calcified (75%) and low-attenuation, indicating early and vulnerable disease. That our severe MetS group shows similar patterns supports the notion that MetS fosters early phenotypes of high-risk coronary atherosclerosis even before symptoms appear. (25)

The observed correlations between metabolic parameters and plaque burden (fasting glucose, triglycerides, waist circumference, BMI, and HDL) align with the mechanistic understanding of MetS: insulin resistance, visceral adiposity, dyslipidaemia, hypertension and low-grade inflammation all promote endothelial dysfunction, vascular remodelling and plaque formation. Several prior studies link number of MetS components with increasing CACS and plaque burden; for example, Kim et al. found a graded increase in CAD progression with number of MetS components. Our data replicate this graded relationship and show that the combined MetS severity score correlates more strongly ($\rho = 0.55$) than single parameters. (26)

A strength of our study is that it quantifies multiple dimensions of subclinical coronary disease (plaque types, burden, high-risk features) in a well-defined high-risk MetS cohort, and shows a robust dose-response relationship with clinical severity of metabolic derangement. It thereby adds value beyond prior studies which either reported large asymptomatic cohorts without MetS severity stratification or progression studies without full burden characterization.

Limitation

First, this is a cross-sectional design; hence, causal inferences or prognostic outcomes cannot be proven. Long-term follow-up would be needed to assess whether the high prevalence and burden of plaque observed translates into future cardiovascular events. Second, the sample size of 100, though adequate for correlations, is modest compared to large screening cohorts, and may limit generalisability. Third, radiation exposure, contrast nephropathy risk and cost limit routine use of CCTA as a screening tool; hence, the value of imaging must be balanced with risk-benefit and cost-effectiveness considerations. Fourth, despite stratification by severity, residual confounding from other risk factors (e.g., smoking, family history) cannot be excluded.

Conclusion:

In conclusion, our study demonstrates that in high-risk MetS patients, subclinical coronary atherosclerosis is highly prevalent, and its burden and high-risk features increase significantly with metabolic severity. Combined with published evidence, these data suggest that CCTA may serve as a powerful imaging adjunct for early detection and risk stratification in MetS, enabling more targeted preventive interventions. Future prospective studies are warranted to assess whether CCTA-guided management translates into improved cardiovascular outcomes in this population.

References:

- Mendis S, Puska P, Norrving B. Global atlas on cardiovascular disease prevention and control. Geneva: World Health Organization; 2011.
- World Health Organisation. Disease and injury country estimates. Global burden of disease. WHO; 2012.
- Finegold JA, Asaria P, Francis DP. Mortality from ischaemic heart disease by country, region, and age. *Int J Cardiol*. 2013;168(2):934–45.
- Resnick HE, Lindsay RS, McDermott MM, Devereux RB, Jones KL, Fabsitz RR, et al. High and low ankle-brachial index and cardiovascular mortality. *Circulation*. 2004;109(6):733–9.
- Heald CL, Fowkes FG, Murray GD, Price JF. Risk associated with ankle-brachial index: systematic review. *Atherosclerosis*. 2006;189(1):61–9.
- Howard BV, Wylie-Rosett J. Sugar and cardiovascular disease. *Circulation*. 2002;106(4):523–7.
- Bots ML, Hoes AW, Koudstaal PJ, Hofman A, Grobbee DE. Carotid intima-media thickness and risk of MI and stroke. *Circulation*. 1997;96(5):1432–7.
- von Scholten BJ, Reinhard H, Hansen TW, Lindhardt M, Petersen CL, Wiinberg N, et al. NT-proBNP and CAC in diabetes. *Cardiovasc Diabetol*. 2015;14:59.
- Greenland P, LaBree L, Azen SP, Doherty TM, Detrano RC. CAC and Framingham score for risk prediction. *JAMA*. 2004;291(2):210–5.
- Polonsky TS, McClelland RL, Jorgensen NW, Bild DE, Burke GL, Guerci AD, et al. CAC score and risk classification. *JAMA*. 2010;303(16):1610–6.
- Bonow RO. Coronary calcium screening—clinical practice. *N Engl J Med*. 2009;361(10):990–7.
- Greenland P, Bonow RO, Brundage BH, Budoff MJ, Eisenberg MJ, Grundy SM, et al. ACCF/AHA consensus on CAC scoring. *J Am Coll Cardiol*. 2007;49(3):378–402.
- Greenland P, Alpert JS, Beller GA, Benjamin EJ, Budoff MJ, Fayad ZA, et al. 2010 ACCF/AHA guidelines for risk assessment. *J Am Coll Cardiol*. 2010;56(25):e50–103.
- Budoff MJ, Hokanson JE, Nasir K, Shaw LJ, Kinney GL, Chow D, et al. CAC progression predicts mortality. *JACC Cardiovasc Imaging*. 2010;3(12):1229–36.
- Raggi P, Callister TQ, Shaw LJ. CAC progression and MI risk. *Arterioscler Thromb Vasc Biol*. 2004;24(7):1272–7.
- McEvoy JW, Blaha MJ, Defilippis AP, Budoff MJ, Nasir K, Blumenthal RS, et al. CAC progression—clinical review. *J Am Coll Cardiol*. 2010;56(20):1613–22.
- Leber AW, Knez A, Becker A, Becker C, von Ziegler F, Nikolaou K, et al. Accuracy of MDCT for plaque composition vs IVUS. *J Am Coll Cardiol*. 2004;43(7):1241–7.
- Achenbach S, Moselewski F, Ropers D, Ferencik M, Hoffmann U, MacNeill B, et al. Detection of non-calcified plaque by MDCT. *Circulation*. 2004;109(1):14–7.
- Mowatt G, Cummins E, Waugh N, Walker S, Cook J, Jia X, et al. Effectiveness of 64-slice CT angiography. *Health Technol Assess*. 2008;12(17):1–143.
- Lim S, Shin H, Lee Y, Yoon JW, Kang SM, Choi SH, et al. MetS and plaque characteristics on 64-slice CT. *Radiology*. 2011;261(2):437–45.
- Wilson PW, Kannel WB, Silbershatz H, D'Agostino RB. Clustering of metabolic factors and CHD risk. *Arch Intern Med*. 1999;159(10):1104–9.
- Isomaa B, Almgren P, Tuomi T, Forsen B, Lahti K, Nissen M, et al. Cardiovascular morbidity in metabolic syndrome. *Diabetes Care*. 2001;24(4):683–9.
- Park GM, An H, Lee SW, Cho YR, Gil EH, Her SH, Park HW, Ahn JM, Park DW, Kang SJ, Kim YH, Lee CW, Yang DH, Kang JW, Lim TH, Kim HK, Choe J, Park SW, Park SJ. Impact of Metabolic Syndrome on Subclinical Atherosclerosis in Asymptomatic Individuals. *Circ J*. 2015;79(8):1799–806. doi: 10.1253/circj.CJ-14-1197. Epub 2015 May 8. PMID: 25959307.
- Kim, L.K., Yoon, J.W., Lee, D.H. et al. Impact of metabolic syndrome on the progression of coronary calcium and of coronary artery disease assessed by repeated cardiac computed tomography scans. *Cardiovasc Diabetol* 15, 92 (2016). <https://doi.org/10.1186/s12933-016-0404-7>